





SOLUTION OF WORKSHEET-17

SUBJECT - STATISTICS

<u>Term</u> : 1st										
Topic	- CENT	TRAL TENDEN		Class: XI						
Full Marks: 15				Date:09 .07. 2020						
Q1.	Select the correct alternative of the following questions.									
	(i) The marks of 5 students in a class test are 1, 2, 4, 7, 8, 11. The median is great than									
		(a) 2	(b)5.5	(c)8	(d) none of these					
	(ii)	The median is (a) all values	used when the set of (b) outlier	observations has (c)equal values	(d) none of these					
	(iii) The marks of 5 students in a class test are 2, 4, 7, 8, 23. The median									
		(a) 2	(b)7	(c)11	(d) none of these					
	(iv)	If all the Obser	then the median is equ	dian is equal to						
			1	(c)-5	(d) none of these					
	(v) Median of -(2n+3),, -1, 0, 1,, (2n+3) is									
		(a) -1	(b) 0	(c) $\frac{n-1}{2}$	(d) ∞					
	(vi) Median of religion of several people									
		(a)n-1	(b) 0	(c) $\frac{n-1}{2}$	(d) none of these					

(vii)	The median can always be calculated of a set having observation							
	(a) countably	infinite	(b	(b) uncountably infinite				
	(c) uncountab	ly finite	(d	(d) none of these				
(viii)	If 5x=9y then and median of x is 7, then median of y is							
	(a) 0	(b)1	(c)0. 5	(d) none of these				
(ix)	The combined harmonic mean depends upon the mean of							
	(a) 1 st set	(b) 2 nd set	(c) both	(d)none of these				
(x)	The composite median is less than the harmonic mean of the given sets which is (a) maximum (b) minimum (c) both (d) none of these							
(xi)	The composite arithmetic mean is greater than the median of the given s which is							
	(a) maximum	(b) minimum	(c) both	(d) none of these				
(xii)	The sum of differences of median from to all the observations except one value is							
	(a) -1	(b) 1	(c) 0	(d) none of these				
(xiii)	Theres are 10 observations with median 3. If 0.3 is added to all the observations then the median of the new set is							
	(a) 3.3	(b) 10	(c) 30	(d) none of these				
(xiv)	There are 10 observations with median 4. If all the observations be added by 4 then of the reciprocal of median of those are							
	(a)0	(b) 2	(c) 1	(d) none of these				
(xv)	The suitable measure to find the central value when all the observations are equal							
	(a) AM	(b) GM	(c) all	(d) none of these				

Prepared by Sanjay Bhattacharya