



**ST. LAWRENCE HIGH SCHOOL**  
**A JESUIT CHRISTIAN MINORITY INSTITUTION**



**WORK SHEET 5**

**Subject : PHYSICS**

CLASS : XII

Date : 8.5.20


Chapter : Electrostatics

Topic : Torque on Dipole, Potential, Potential due to  
Dipole on Axial and Perpendicular bisector point.

**Multiple Choice Question :**

**1 x 15 = 15**

1. A dipole of dipole moment  $\vec{p}$  is placed in uniform electric field  $\vec{E}$  then torque acting on it is given by :  
(a)  $\vec{\tau} = \vec{p} \cdot \vec{E}$  (b)  $\vec{\tau} = \vec{p} \times \vec{E}$   
(c)  $\vec{\tau} = \vec{p} + \vec{E}$  (d)  $\vec{\tau} = \vec{p} - \vec{E}$
2. An electric dipole is placed at an angle of  $30^\circ$  with an electric field intensity  $2 \times 10^5 \text{ N C}^{-1}$ . It experiences a torque equal to 4 N m. The charge on the dipole, if the dipole length is 2 cm, is  
(a) 8 mC (b) 2 mC (c) 5 mC (d) 7  $\mu\text{C}$
3. An electric dipole with dipole moment  $\vec{p} = (2\hat{i} + 3\hat{j}) \text{ cm}$  is kept in electric field  $\vec{E} = 4\hat{i} \text{ N/C}$ . The torque acting on it is :  
(a)  $-12\hat{k} \text{ (Nm)}$  (b)  $8\hat{k} \text{ (Nm)}$  (c)  $12\hat{k} \text{ (Nm)}$  (d)  $-8\hat{k} \text{ (Nm)}$
4. Number of Statvolt corresponding to 1 volt is :  
(a)  $\frac{1}{100}$  (b)  $10^9$  (c)  $\frac{1}{300}$  (d) 300
5. Four point charges each  $+q$  is placed on the circumference of a circle of diameter  $2d$  in such a way that they form a square. The potential at the centre of the circle (in CGS) is :  
(a) 0 (b)  $\frac{4q}{d}$  (c)  $\frac{4d}{q}$  (d)  $\frac{q}{4d}$
6. The radius of a soap bubble whose potential is 16 V is doubled. The new potential of the bubble is :  
(a) 2 V (b) 4 V (c) 8 V (d) 16 V
7. Four electric charges  $+q, +q, -q$  are placed at the corners of a square of side  $2L$ . The electric potential at point A midway between the two charges  $+q$  and  $+q$  is :



- (a)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} (1 + \sqrt{5})$  (b)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} \left(1 + \frac{1}{\sqrt{5}}\right)$  (c)  $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} \left(1 - \frac{1}{\sqrt{5}}\right)$  (d) zero

8. The electric potential at a point on the axis of an electric dipole depends on the distance  $x$  of the point from the dipole as :
- (a)  $\propto x$  (b)  $\propto \frac{1}{x}$  (c)  $\propto \frac{1}{x^2}$  (d)  $\propto \frac{1}{x^3}$
9. The electric potential at the surface of an atomic nucleus ( $Z = 50$ ) of radius of  $9 \times 10^{-15} \text{ m}$  :
- (a) 80V (b) 9V (c)  $9 \times 10^5 \text{V}$  (d)  $8 \times 10^5 \text{V}$
10. Two charges  $-10\text{C}$  and  $+10\text{C}$  are placed 10 Cm apart. Potential at the centre of the line joining the two charges is
- (a) zero (b) 2 V (c)  $-2 \text{ V}$  (d) None of these
11. Two charges each equal to  $q$  are placed at the corners of a square of side  $l$ . The electric potential at the centre of the square is :
- (a)  $\frac{1}{4\pi\epsilon_0} \frac{4q}{l}$  (b)  $\frac{1}{4\pi\epsilon_0} \frac{4q}{\sqrt{2}l}$  (c)  $\frac{1}{\pi\epsilon_0} \frac{\sqrt{2}q}{l}$  (d)  $\frac{1}{\pi\epsilon_0} \frac{2q}{l}$
12. Two concentric spheres of radii  $R$  and  $r$  have similar charges with equal surface densities ( $\sigma$ ). What is the electric potential at their common centre?
- (a)  $\frac{\sigma}{\epsilon_0}$  (b)  $\frac{\sigma}{\epsilon_0} (R - r)$  (c)  $\frac{\sigma}{\epsilon_0} (R + r)$  (d) None of the above.
13. At a point A, there is an electric field of 500 V/m and potential of 3000 V. The distance between the point charge and A is :
- (a) 6 m (b) 12 m (c) 36 m (d) 144 m
14. Potential at a point on the perpendicular bisector of a dipole is :
- (a) zero (b) 1 (c)  $\frac{q}{2l}$  (d)  $\frac{l}{q}$
15. Potential at a distance  $r$  from the mid point of a dipole of length  $2l$  on the axis of it is :
- (a) zero (b)  $\frac{1}{4\pi\epsilon_0} \cdot \frac{p}{r^2 - l^2}$  (c)  $\frac{1}{4\pi\epsilon_0} \cdot \frac{2p}{r^2 - l^2}$  (d)  $\frac{P}{r^2 - l^2}$

***Ambarnath Banerjee***