

ST. LAWRENCE HIGH SCHOOL



A JESUIT CHRISTIAN MINORITY INSTITUTION

Sub: Biological Sciences Class: XIDate: 25.07.2020

Animal KIngdom: Porifera, Cnidaria and Ctenophora (Part-I) F.M:15

		WORKSHEET	<u> – 27</u>	(1x15=15)
i)	Outer cell layer of Poriferans is called			
(1)	Ectoderm	(2)Pinacoderm	(3)Choanoderm	(4) Endoderm
ii)	Which of the following layers are present in between two cell layers in the body of			
	Poriferans?			
(1)	Epidermis	(2) Mesenchyme	(3) Choanoderm	(4)Coelom
iii)	The pores that help in the entry of the water is			
(1)	Ostia	(2) Osculum	(3) Nematocysts	(4) None of these
iv)	Spongocoel remain lined by collared cells called			
(1)	Choanocytes	(2) Amoebocyte	(3) Pinacocytes (4) Scleroblasts	
v)	Which of the following is a fresh water sponge?			
(1)	Sycon	(2)Spongilla	(3)Euplectella	(4)Euspongia
vi)	Which of the following organisms is radially symmetric?			
(1)	Hydra	(2)Aurelia	(3)Obelia	(4) All of these
vii)	The specialised cells which is present in the ectoderm of Cnidaria are called-			
(1)	Pinacocytes	(2) Cnioblasts	(3) Amoebocytes	(4)Scleroblasts
viii	The opening of gastrovascu	lar cavity that maintai	n connection with the	cavity in Cniodarians
	is -			
(1)	Ostia	(2)Osculum	(3)Hypostome (4) No	ne of these
ix)	The asexual phase in the life cycle of Cnidarians is called-			
(1)	Polyp	(2)Medusa	(3)Both (1) and (2)	(4)None of these
x)	The number of anal pores present in Ctenophora is -			
(1)	1	(2)2	(3) 3	(4) 4
xi)	The number of ciliary comb plates is-			
(1)	1	(2) 3	(3) 5	(4) 8
xii)	Statolith is-			
(1)	Respiratory Organ	(2)Circulatory Organ	(3)Sense Organ	(4)Nervous System
xiii	Colloblast cells or Lasso Cel	ls are present in-		
((1) Cnidaria	(2)Ctenophora	(3)Porifera	(4)All of these
xiv	Stomodeum is found in			
((1) Hydra	(2) Obelia	(3)Aurelia	(4)Pleurobrachia
xv)	The similarity between <i>Cter</i>	noplana and Obelia is		
((1) Presence of Cnidoblast	(2) Alternation of ger	neration (3)Bira	adial Symmetry
	(4) Nerve net			
