

A JESUIT CHRISTIAN MINORITY INSTITUTION

Sub: Biological Sciences Class: XIDate: 8.08.2020

FOR GOD AND COUNTRY



| <u>Animal Kingdom: Chordata (Part-I)</u> | | | F.M:15 |
|--|----------------------|----------------------|---------------------|
| | WORKSHEET | <u>- 30</u> | (1x15=15) |
| i) Which of the following is not an identifying feature of chordates? | | | |
| (1) Notochord | (2)Dorsal nerve chor | d (3) Pharyngea | l gill slits |
| (4) True coelom | | | |
| ii) Which of the following does not have an exoskeleton? | | | |
| (1) Cockroach | (2) Butterfly | (3) Balanoglossus | (4) Prawn |
| iii) The dorsal nerve cord is sol | id in | | |
| (1) Hemichordata | (2) Urochordata | (3) Cephalochordata | (4) Vertebrata |
| iv) Which of the following is absent in Hemichordata? | | | |
| (1) Pharyngeal gill slits | (2) Nerve cord | (3) stomochord | (4) Post -anal tail |
| v) Notochord is absent in- | | | |
| (1) Branchiostoma | (2) Asymmetron | (3)Saccoglossus | (4) All of |
| these | | | |
| vi) Retrogressive metamorphosis is observed in | | | |
| (1) Urochordata | (2)Hemichordata | (3)Cephalochordata | (4) Vertebrata |
| vii) In which of the following organisms does the adult body lack notochord? | | | |
| (1) Doliolum | (2) Branchiostoma | (3) Asymmetron | (4)Petromyzon |
| viii)Which of the following structure is found in cephalochordates? | | | |
| (1) Cilia | (2) Tentacles | (3)Flagella | (4) Pili |
| ix) In Cephalochordata excretion occurs by | | | |
| (1) Kidney | (2)Protonephridia | (3)Nephridia | (4)None of these |
| x) Which of the following has a fish-like structure? | | | |
| (1) Branchiostoma | (2)Asymmetron | (3) Both (1) and (2) | (4) None of these |
| xi) Brain is a swelling of | | | |
| (1) Nerve cord | (2) Notochord | (3) Appendages | (4) Vertebral |
| column | | | |
| xii) The excretory organ in vertebrates is | | | |
| (1) Kidney | (2) Protonephridia | (3)Nephridia | (4) Green glands |
| xiii) Which of the following has a cranium? | | | |
| (1) Hemichordata | (2)Cephalochordata | (3) Urochordata | (4)Vertebrata |
| xiv)The number of cranial nerves present in vertebarates is | | | pairs |
| (1) 2-4 | (2) 4-8 | (3) 8-10 | (4) 10-12 |
| xv) In hemichordates, the body is divisible into | | | |
| (1) Proboscis | (2) Collar | (3) Trunk | (4) All of these |
| | - | - | - |

Manjaree Guha