



# ST. LAWRENCE HIGH SCHOOL

A JESUIT CHRISTIAN MINORITY INSTITUTION

## WORKSHEET-16(CLASS-11)

### TOPIC- STRUCTURE OF ATOM

### SUBTOPIC- QUANTUM NUMBER



SUBJECT – CHEMISTRY

DURATION – 30 mins

F.M. - 15

DATE -02.07.20

1.1 How many orbitals can have the following set of quantum number(s):

$$n = 3, l = 1, m_l = 0$$

a) 3 b) 1 c) 4 d) 2

1.2 Electronic configuration of the outer shell of the element Gd with atomic number 64 is:

(a)  $4f^4 5d^5 6s^1$  (b)  $4f^3 5d^5 6s^2$  (c)  $4f^5 5d^4 6s^1$  (d)  $4f^7 5d^1 6s^2$

1.3 Maximum number of electrons in a subshell can be:

(a)  $4l + 2$  (b)  $4l - 2$  (c)  $2n^2$  (d)  $2l + 1$

1.4 The orientation of atomic orbitals depends on their:

(a) Spin quantum number (b) magnetic quantum number (c) azimuthal quantum number  
(d) Principal quantum number

1.5 Number of unpaired electrons in  $N^{2+}$ :

(a) 3 (b) 1 (c) 2 (d) 0

1.6 If the electronic structure of oxygen atom is written as  $1s^2 2s^2 2p^4$  it would violate:

(a) Hund's rule (b) Pauli's exclusion principle (c) Both Hund's and Pauli's principles  
(d) None of these

1.7 Which quantum number(s) do 2s and 2p orbitals have in common?

(a) n and l (b) n (c) l and  $m_l$  (d) l

1.8 For a principal quantum number n, how many atomic orbitals are possible?

a)  $n^2$  b)  $n + 1$  c)  $2n$  d) n

1.9 Which set of quantum numbers uniquely defines one of the electrons in an atomic orbital with  $n = 2$  and  $l = 0$ ?

(a)  $n = 2, l = 0, m_l = 1, m_s = +\frac{1}{2}$  b)  $n = 2, l = 0, m_l = 0, m_s = +\frac{1}{2}$   
(c)  $n = 2, l = 0, m_l = 1, m_s = +1$  d)  $n = 2, l = 0, m_l = 0, m_s = +1$

**1.10 Element Z has the ground state electronic configuration  $1s^2 2s^2 2p^3$ . In which group does it belong?**

(a) 5 (b) 15 (c) 3 (d) 13

**1.11 A set of orbitals for which the quantum number  $l = 2$  is:**

(a) 7-fold degenerate (b) non-degenerate (c) 3-fold degenerate (d) 5-fold degenerate

**1.12 For the principal quantum number  $n = 4$ , it is possible to have:**

(a) only s and p orbitals (b) only s, p, d and f orbitals (c) only an s orbital (d) only s, p and d orbitals

**1.13 “No two electrons in atom can have the same set of all four quantum numbers”-**

(a) Pauli’s exclusion principle (b) Hund’s rule of maximum spin multiplicity  
(c) The Aufbau Principle (d) None of these

**1.14 The electronic configuration of Cr is:**

a)  $[Ar]4s^2 3d^4$  (b)  $[Ar]4s^1 3d^4$  (c)  $[Ar]3d^6$  (d)  $[Ar]4s^1 3d^5$

**1.15 Find the number of unpaired electrons present in  $Fe^{2+}$ :**

(a) 6 (b) 4 (c) 5 (d) 3

**PREPARED BY: MR. ARNAB PAUL CHOWDHURY**